

CROSSCUT SELECT™

Post-emergent brush, woody plant and broadleaf weed control

ACTIVE INGREDIENTS

Triclopyr – Group 4
2,4-D – Group 4

% ACTIVE INGREDIENTS

Triclopyr BEE 16.5%
2,4-D 34.4%

FORMULATION

Liquid

CHEMICAL FAMILY

Chlorinated
phenoxy-systemic herbicide

BRAND ALTERNATIVE

Crossbow®

MODE OF ACTION

Growth regulator, induces rapid, uncontrolled growth until death

RESTRICTED USE

No

SIGNAL WORD


Caution

PACKAGING

12 x 1 qt, 4 x 1 gal
2 x 2.5 gal

LABELED USES

Commercial and recreational turf
Commercial sod
Rangeland
Permanent grass pastures
Conservation reserve program (CRP) acres
Fence rows
Non-irrigation ditchbanks
Industrial sites



CrossCut Select™ is a post-emergent herbicide that provides control of most woody plants and brush, as well as annual and perennial broadleaf weeds. Formulated as a low volatile ester, the combination of triclopyr and 2,4-D provides outstanding weed control in both cool and warm weather.

PRODUCT BENEFITS

- Provides excellent control of brush and woody plants
- Controls broadleaf weeds with no harm to turf
- Low volatility formulation
- Can be applied by all types of sprayers
- Rainfast within 2 hours post application
- Flexible application timing – can be applied in-season or during dormant periods
- Apply to actively growing foliage for best results

FOR USE ON TURF

Broadcast treatment of commercial and recreational turf, commercial sod farms

To be applied only under the direct supervision of Commercial applicators responsible for turf weed control programs.

Apply 2 to 4 pints (.5 to 1 lb. 2,4-D ai) of **CrossCut Select™** in enough water to make 20 to 200 gallons total spray per acre to control broadleaf weeds growing in tall fescue, bluegrass, or perennial ryegrass turf. Do not use on other grass species, such as bentgrass or St. Augustine grass, unless injury can be tolerated.

Restrictions

For broadcast application on commercial, and recreational turf

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

Post-emergence

- Limited to 2 applications per year.
- Use a maximum of 4 pints **CrossCut Select™** (1.5 lbs. 2,4-D ai) per acre per application.
- The maximum seasonal rate is 8 pints **CrossCut Select™** per acre (2.0 lbs. 2,4-D ai/A), excluding spot treatments.



Prime Source, a Division of Albaugh, LLC

1525 NE 36th Street, Ankeny, IA 50021 • www.primesource-albaugh.com

©2021 Albaugh, LLC. Prime Source and Select are trademarks of Albaugh, LLC. Crossbow is a registered trademark of Dow AgroScience LLC. Always read and follow label instructions.

CROSSCUT SELECT™

Post-emergent brush, woody plant and broadleaf weed control

FOR USE ON RANGELAND AND PASTURE

Broadcast treatment (ground equipment and helicopter)

Apply up to 4 quarts of **CrossCut Select™** per acre in sufficient water to deliver 10 to 30 gallons of total spray per acre. Actively growing weeds are most susceptible. Optimal application time for biennial and winter annual weeds is the rosette stage. Hard to control weeds such as field bindweed, chicory, dogfennel, goldenrod, horsenettle, kudzu, milkweed, perennial sowthistle, leafy spurge, and Canada thistle may require retreatment for complete control.

Spot treatment

To control broadleaf weeds in small areas with a hand sprayer, mix 4 to 6 fl. oz. of **CrossCut Select™** in 3 gallons of water. Spray to thoroughly wet all foliage.

Restrictions

For rangeland and pasture

- Do not apply more than 4 quarts per acre per application.
- Do not make more than one application per year.
- Pre-harvest interval is 14 days
- Except for lactating dairy animals there are no grazing restrictions
- Do not apply more than 1 lb. 2,4-D ai per application (2 qts. product) for susceptible annual and biennial broadleaf weeds.

Weeds controlled

| | | | |
|----------------|--------------------------------------|----------------------------------|-------------|
| Black medic | Chickweeds (common and mouse ear) | Field Bindweed | Poison ivy |
| Bull thistle | Cinquefoil | Ground ivy | Poison oak |
| Burdock | Clovers | Henbit | Speedwell |
| Buttercup | Creeping beggar weed | Lambsquarters | Smartweed |
| Blackberry | Dandelion | Oxalis (stricta and corniculata) | Vetch |
| Canada thistle | Docks | Plantain | Wild violet |

Woody plant control (easy to control species)

6 quarts/acre broadcast application or 1 to 1.5% mixtures for high-volume foliar applications.

| | | | |
|-------------------------------------|--------------|-------------------------|----------------------|
| Alder | Dogwood | Sumac | Speedwell |
| Ash | Elderberry | Sycamore | Smartweed |
| Beech | Hawthorn | Tamarack | Vetch |
| Blackberry cherry (except black) | Poison ivy | Wax myrtle (top growth) | Wild violet and more |
| Cottonwood | Poison oak | White oak | |
| | Scotch broom | Wild grape | |

Woody plant control (harder to control species)

Apply high-volume applications, 1.5% mixture, conventional basal or dormant stem applications. A broadcast rate of 8 quarts/A will increase the degree of control of these species.

| | | |
|---|---------------------------|-------------------------------|
| Buckbush (<i>Symphoricarpos</i> spp.) (suppression) | Hazel | Salmonberry (suppression) |
| Common persimmon (suppression) | Honeylocust (suppression) | Sweetgum |
| Elm (except winged elm) | Pine (suppression) | Trumpet creeper (suppression) |
| | Russian olive | Virginia creeper |