

REGULATE SELECT™

Systemic fungicide for disease control

ACTIVE INGREDIENT

Metalaxyl – Group 4

% ACTIVE INGREDIENT

Metalaxyl 44.08%

FORMULATION

Flowable

CHEMICAL FAMILY

Phenylamide fungicide

BRAND ALTERNATIVE

Subdue MAXX®

MODE OF ACTION

Systemic

RESTRICTED USE

No

SIGNAL WORD

Caution

PACKAGING

4 x 1 gal

LABELED USES

- Golf courses
- Lawns
- Residential landscape areas
- Ornamentals
- Instructional landscape areas
- Public landscape areas
- Commercial landscape areas
- Industrial landscape areas
- Parks
- Recreational areas
- Athletic fields
- Sod farms



Regulate Select™ is a systemic fungicide that provides control of certain diseases in turf, ornamentals & greenhouses. With a unique water-based formulation, **Regulate Select™** is low odor and mixes thoroughly and easily into most commonly used fungicide and fertilizer tank mixes. **Regulate Select™** works both through contact action with the root system, as well as systemic movement through the entire plant.

PRODUCT BENEFITS

- Effectively controls *Pythium* in all turf species
- Low odor
- Unique water-based flowable formulation
- Exceptional tank mix compatibility
- Same use rate as Subdue Maxx®
- For use on both established and newly seeded turf



Prime Source, a Division of Albaugh, LLC
1525 NE 36th Street, Ankeny, IA 50021 • www.primesource-albaugh.com

©2021 Albaugh, LLC. Prime Source and Select are trademarks of Albaugh, LLC. Subdue MAXX is a registered trademark of Syngenta Group Company. Always read and follow label instructions.

REGULATE SELECT™

Systemic fungicide for disease control

APPLICATION INFORMATION

Ornamentals

Use this product on container, bench, or bed-grown ornamentals in greenhouses or outdoor nurseries, and for use on ornamentals grown for indoor and outdoor landscaping, for control of damping-off and root and stem rot diseases caused by *Pythium* and *Phytophthora*.

This product may be applied through irrigation systems, as a soil drench or as a soil surface spray, or incorporated into a pre-potting growing media for subsequent seeding or transplanting of ornamentals.

Citrus in nurseries and landscape plantings (non-bearing)

Use this product on non-bearing citrus for control of citrus foot rot, root rot, and trunk canker caused by *Phytophthora* spp. Apply to the soil as a drench or as a spray in a banded application. Make the first application of this product at the time of planting. Make repeat applications at 3-month intervals during the period when trees are actively growing.

Conifers in nurseries and plantations (including Christmas trees)

Use of this product will aid in the control of *Phytophthora* root rot when used in conjunction with good cultural practices. The use of this product will not overcome poor management practices, such as planting on sites that are prone to flooding or are poorly drained. This product will not revitalize trees showing moderate to severe disease symptoms.

Turfgrass

This product controls *Pythium* blight and *Pythium* damping-off in turf, yellow tuft (downy mildew) in bluegrass, and downy mildew in St. Augustine grass. For optimum results, begin applications before symptoms occur when conditions first favor disease. During periods of prolonged conditions favorable for disease development, use on a 14-day schedule. For long-term control of *Pythium* in areas when using seed treated with the active ingredient contained in this product, make an application of this product 7 to 10 days after seeding. For control of other diseases of turf, use **Propiconazole Select™** alone or in a tank-mix combination with this product. Refer to the **Propiconazole Select™** label for rates, precautions, restrictions, etc.

Labeled diseases

Pythium

Phytophthora spp.

Downy mildew

Yellow tuft

