

CARFENTRAZONE-ETHYL GROUP 14 HERBICIDE



RightLine®

ADDIT™ HERBICIDE

INTENDED FOR COMMERCIAL USE

ACTIVE INGREDIENT:	By Wt.
Carfentrazone-ethyl.....	21.58%
OTHER INGREDIENTS:	78.42%
TOTAL:	100.00%

This product contains 1.9 pounds active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See inside label booklet for additional Precautionary Statements, and Directions for Use including Storage and Disposal instructions.

EPA Reg. No.: 95009-2-93051

EPA Est. No.: 102434-GA-1 (AC), 89332-GA-001 (MA),
86869-NC-001 (SM)

(Letter(s) in the lot number correspond to letter(s)
following the EPA Est No.)



1 97644 63342 5

Net Contents: 0.5 gallon (64 fl. oz.)

Distributed By:
RightLine, LLC
950 Falcon Drive
Malden, MO 63863

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- **DO NOT** induce vomiting unless told to do so by a poison control center or doctor.
- **DO NOT** give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12 PM Pacific Standard Time. In the event of a medical emergency, call your poison control center at 1-800-222-1222.

**For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof chemical-resistant gloves, and shoes plus socks.

User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling the product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Carfentrazone-ethyl is very toxic to algae and moderately toxic to fish. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the high-water mark, except as specified on this label. **DO NOT** contaminate water when disposing of equipment washwater.

Fish Advisory Statement:

This product may be hazardous to aquatic organisms, particularly in clear, shallow water bodies that are adjacent to treated areas. Transport to water by runoff or spray drift of this product in areas where surface water is present, or intertidal areas below the mean high water mark, should be avoided. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

For ground water:

Residues of this chemical have properties and characteristics associated with chemicals detected in ground water. Residues of this chemical may leach into ground water if the chemical is used in areas where soils are permeable, particularly where the water table is shallow.

For Surface Water:

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of carfentrazone-ethyl residues from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Non-target Organism Advisory Statement:

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by minimizing spray drift.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT mix or allow coming into contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Re-entry Statement: DO NOT allow people (other than applicator) or pets on treatment area during application. **DO NOT** enter treatment area until spray has dried.

WEED RESISTANCE MANAGEMENT

For resistance management, RightLine ADDIT Herbicide is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to RightLine ADDIT Herbicide and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of RightLine ADDIT Herbicide or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact RightLine, LLC retailer or representative.

Report any incidence of non-performance of this product against a particular weed species to your RightLine, LLC retailer or representative. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemical means to remove escapes, as practical, with the goal of preventing further seed production.

PRODUCT INFORMATION

RightLine ADDIT Herbicide is emulsion oil in water formulation. RightLine ADDIT Herbicide is to be mixed with water, liquid fertilizer or mixtures of water and liquid fertilizer and adjuvants and applied to non-crop areas including turfgrass for selective postemergence control of broadleaf weeds

Weed control is optimized when the product is applied to actively growing weeds. RightLine ADDIT Herbicide is a contact herbicide. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation.

Extremes in environmental conditions including temperature, moisture, soil conditions, and cultural practices may affect the activity of RightLine ADDIT Herbicide, symptoms may be accelerated under moist conditions. Weed control may be reduced when weeds are hardened off by drought and become less susceptible to RightLine ADDIT Herbicide.

RightLine ADDIT Herbicide is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications are advised not to be made within 6 to 8 hours of either rain or irrigation or when heavy dew is present on the crop. Environmental conditions and with certain spray tank additives may increase herbicidal symptoms on the crop.

TANK MIXTURES

RightLine ADDIT Herbicide may be tank-mixed with other registered herbicides for controlling broader spectrum weeds. Refer to this and other product's labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank mix partner. When preparing a new tank mix conduct an appropriate compatibility test by mixing proportional amounts of all spray ingredients in a test vessel (jar) prior to tank mixing with other products. Shake the mixture vigorously and allow it to stand for five to ten minutes. Rapid precipitation of the ingredients and failure to re-suspend when shaken indicates that the mixture is incompatible and must not be applied. Provided the jar test indicates the mixture to be compatible, prepare the tank mixture as follows: Fill the tank one fourth full with water. With the agitator operating, add the labeled amounts of ingredients using the following order: dry granules first and liquid suspensions (flowables) second. As the agitation continues and the tank is filled with water add emulsifiable concentrate products third followed by the addition of water-soluble products.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

ADJUVANT USE REQUIREMENTS

The use of a quality spray adjuvant is required for optimum performance. Refer to the individual sections of this label for specific adjuvant type and use rates.

MIXING INFORMATION

Mixing and Loading Instructions

Start by filling the tank with $\frac{3}{4}$ of the desired volume of clean water and, with agitation, add the proper amount of RightLine ADDIT Herbicide. Complete filling the spray tank to the desired volume. Maintain sufficient agitation to keep materials in solution during both mixing and application and until the spray tank has been emptied. For tank mixtures, follow your local extension guidelines for mixing order. General guidelines are: add dry materials first and agitate until mixed; then EW or water soluble liquids; then EC formulations; then, add adjuvants last. Ensure the compatibility of other products and/or liquid fertilizers with RightLine ADDIT Herbicide before mixing them together in the spray tank.

Mixing Precautions

Avoid the overnight storage of RightLine ADDIT Herbicide spray mixtures. If spray solution is stored overnight or longer, thoroughly agitate spray mixture before applying the solution.

Mixing Restrictions

- **DO NOT** premix RightLine ADDIT Herbicide spray solutions in nurse tanks. Maintain continuous and adequate spray solution agitation until all the spray solution has been used.
- **DO NOT** use with tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer spray solution to alter the pH range as appropriate.

SPRAY EQUIPMENT CLEAN-OUT

Many new pesticides are very active at low rates, especially to sensitive crops. Residues left in mixing equipment, spray tanks, hoses, spray booms and nozzles can cause crop effects if they are not properly cleaned. As soon as possible after spraying RightLine ADDIT Herbicide and before using the sprayer equipment for any other applications, the sprayer equipment must be thoroughly cleaned using the following procedure. In addition, users must take appropriate steps to ensure proper equipment clean-out for any other products mixed with RightLine ADDIT Herbicide as required on the other product labels. More complete cleaning can be achieved if the spray system is cleaned immediately following the application.

1. Drain sprayer tank, hoses, spray boom and spray nozzles. Use a high-pressure detergent wash to remove physical sediment and residues from the inside of the sprayer tank and thoroughly rinse. Then, thoroughly flush sprayer hoses, spray boom and spray nozzles with a clean water rinse. Remove and clean spray tips and all filters and screens (tank, spray hose and spray tips) separately in the ammonia solution of Step 2.

2. Next, prepare a sprayer cleaning solution by adding three gallons of ammonia (containing at least 3% active) per 100 gallons of clean water. Prepare sufficient cleaning solution to allow the operation of the spray system for a minimum of 15 minutes to thoroughly flush hoses, spray boom and spray nozzles.

3. Convenient and thorough cleaning of the sprayer can be achieved if the ammonia solution or fresh water is left in the spray tank, hoses, spray booms and spray nozzles overnight or during storage.

4. Before using the sprayer, completely drain the sprayer system. Rinse the tank with clean water and flush through the hoses, spray boom, and spray nozzles with clean water.

5. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State, and local regulations and guidelines.

DO NOT apply sprayer cleaning solutions or rinsate to sensitive crops, lawns, ornamentals, or gardens.

DO NOT store the sprayer overnight or for any extended period of time with RightLine ADDIT Herbicide spray solution remaining in the tank, spray lines, spray boom plumbing, spray nozzles or strainers.

If the sprayer has been stored or idle, purge the spray boom and nozzles with clean water before beginning any application.

Should small quantities of product remain in inadequately cleaned mixing, loading and/or spray equipment, they may be released during subsequent applications potentially causing effects to certain crops and other vegetation. RightLine, LLC accepts no liability for any effects due to inadequately cleaned equipment.

MANDATORY SPRAY DRIFT

Ground Applications

- Apply with the nozzle height specified by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
 - Pressure - Use the lowest spray pressure specified for the nozzle to produce the target spray volume and droplet size.
 - Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.
- **BOOM HEIGHT - Ground Boom**
Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.
 - **SHIELDED SPRAYERS**
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
 - **TEMPERATURE AND HUMIDITY**
When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
 - **TEMPERATURE INVERSIONS**
Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

- **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

- **Boom-less Ground Applications:**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

- **Handheld Technology Applications:**

Take precautions to minimize spray drift.

APPLICATION INSTRUCTIONS

TURF AND ORNAMENTAL

Product Information

RightLine ADDIT Herbicide is most effective when applied to actively growing winter and summer annual seedlings. RightLine ADDIT Herbicide also increases the speed of kill and expands the spectrum of control of mature and perennial broadleaf weeds (See "Weeds Controlled" section) when applied in a tank mixture with other postemergence herbicides.

Extremes in environmental conditions e.g. excessively high or low moisture or temperature may affect the activity of RightLine ADDIT Herbicide. Under warm, moist conditions appearance of herbicide symptoms may be accelerated while under very dry or cool conditions the expression of herbicidal symptoms is delayed. However, RightLine ADDIT Herbicide remains highly effective under both cool and warm environmental conditions. Weeds hardened by drought and/or extremely high temperatures are less susceptible to this product and the higher rates in the range are required under those conditions.

Precautions

Avoid spray drift onto non-target susceptible plants including vegetables, flowers, ornamentals, trees, shrubs and other desirable plants.

Restrictions

DO NOT apply this product to carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present.

DO NOT apply when conditions are conducive to spray drift or poor spray coverage.

(continued)

DO NOT apply more than 6.7 fluid ounces of product (0.10 lbs. a.i. of carfentrazone-ethyl) per acre per application or more than 26.8 fluid ounces (0.40 lbs. a.i. of carfentrazone-ethyl) per acre per season. The minimum retreatment interval for sequential broadcast applications of this product is two (2) weeks.

DO NOT apply this product by aerial application to turf or ornamentals.

DO NOT apply this product through any type of irrigation system.

Turfgrass and Sites

RightLine ADDIT Herbicide is intended to be applied by lawn care operators, maintenance applicators, and service technicians for use in ornamental lawns and turfgrass, as well as stone-covered yards established in institutional, ornamental, and residential/domestic sites. Institutional sites are defined as turf areas and stone-covered yards around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sports facilities, golf courses, and office buildings. Ornamental sites include turfgrass and stone-covered yards established around residences, parks, streets, retail outlets, cemeteries, industrial buildings and institutional buildings. Residential/domestic sites are defined as areas associated with household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

Turfgrass Tolerance

Turfgrasses tolerant to application of RightLine ADDIT Herbicide are listed below. RightLine, LLC does not recommend application to turfgrasses not listed.

Cool Season Grasses (creeping bentgrass, colonial bentgrass, annual bluegrass, Kentucky bluegrass, fine fescues, red fescues, tall fescue, perennial ryegrass)

Established cool season grasses are generally tolerant to applications of RightLine ADDIT Herbicide at labeled rates. Tall fescue may exhibit a slight yellowing discoloration occurring within 3 to 5 days after application under some conditions. Recovery typically occurs within 4 to 7 days. If such injury or any injury cannot be tolerated, apply to a small test area before treating large areas to assure tolerance of the host turf species.

Note

Be aware of and observe all label restrictions regarding turf tolerance of companion products when RightLine ADDIT Herbicide is tank mixed with another product.

Warm Season Grasses (common and hybrid bermudagrass, bahiagrass, buffalograss, centipedegrass, seashore paspalum, St. Augustine grass, zoysiagrass)

Established warm season grasses listed above are generally tolerant to applications of RightLine ADDIT Herbicide but may be susceptible to transitory yellowing when they are under stress. Stress is typically associated with but not limited to extreme high or low temperatures, disease infection or insect infestation, extreme high or low moisture conditions, or transition into or out of dormancy.

Under such conditions injury in the form of transitory yellowing discoloration may occur within 3 to 7 days of application. If such injury or any injury cannot be tolerated, apply RightLine ADDIT Herbicide in the manner you will be using it to a small test area before treating large areas to assure tolerance of the turf.

Note

Be aware of and observe all label restrictions regarding turfgrass tolerance of companion products when RightLine ADDIT Herbicide is tank mixed with another product.

Newly Seeded , Sodded or Sprigged Turfgrass

RightLine ADDIT Herbicide can be applied to the following species of turfgrass at 7 days or more after emergence for species established by both seeding and sprigging: hybrid bermudagrass, common bermudagrass, St. Augustine, creeping bentgrass, tall fescue, perennial ryegrass and Kentucky bluegrass. Application to zoysiagrass must be delayed until at least 14 days after emergence to avoid extended discoloration. Slight discoloration may be observed immediately after application for hybrid bermudagrass, however, normal turf color returns by 12 days after emergence if no other stresses are present. Areas treated with RightLine ADDIT Herbicide may be seeded or overseeded one day following application.

Dormant Turf

Applications to dormant warm-season grasses are permitted. Avoid applications when warm-season grasses are transitioning into or out of dormancy.

Application Directions

Spray Coverage

RightLine ADDIT Herbicide is a contact herbicide with little or no residual activity. It produces herbicidal symptoms only in the portions of the target weeds with which it comes into direct

(continued)

contact. Therefore, to achieve maximum effectiveness of this product, select a spray volume and nozzle system that ensures thorough and uniform coverage and at the same time minimizes the spray fines. Spray droplets larger than 400 microns may reduce coverage and result in loss of weed control. Applications must be made using spray volumes of 20-175 gallons per acre (0.5 to 4.0 gallons per 1000 sq.ft.). Use higher spray volumes when there is a dense weed population or turfgrass canopy or where uniform coverage is difficult to obtain.

Spot Treatments (Applications with hand operated sprayers such as backpack sprayers, compression sprayers, knapsack sprayers)

For spot treating weeds with hand-held or backpack sprayers mix 0.030 fluid ounces (0.9 milliliters) of this product in one to two gallons of water to treat 1000 square feet. To avoid overdosing when using a handheld wand, use a flat fan nozzle and maintain the nozzle at a uniform specified height over the canopy. Avoid back-and-forth and side-to-side motion with the wand. Hold the wand stationary and move forward at a uniform pace over the area to be treated.

Use of Adjuvants

Addition of surface active agents (surfactants) or other adjuvants may improve effectiveness of RightLine ADDIT Herbicide against weeds with difficult-to-wet leaf surfaces or weeds that are mature or hardened by harsh environmental conditions. Adjuvant materials may also reduce the selectivity to some turfgrass varieties and increase the potential for turf injury. Follow all adjuvant manufacturer use guidelines and also read and observe all companion product label statements, precautions and restrictions regarding use of adjuvants when RightLine ADDIT Herbicide is used in tank mixtures with other products.

Weeds Controlled

RightLine ADDIT Herbicide can be used to selectively control weeds anytime during the year except as instructed elsewhere in this label. Application rates for control of weeds range from 0.9 to 2.1 fluid ounces per acre as described below. Use the lower rates in a range for susceptible weed species and for optimal environmental conditions. Use the higher rates in the range for dense infestations of perennial weeds, and for adverse/extreme environmental conditions.

(continued)

Category 1: Used as a stand-alone product

Apply RightLine ADDIT Herbicide at rates of 1.0 to 2.1 fluid ounces of product per acre (0.023 to 0.048 fluid ounces per 1000 square feet) for control of seedling (4 inches tall or less) winter and summer annual weeds listed below:

Annual morningglory spp.	Field pennycress	London rocket
Annual sowthistle	Field speedwell	Nightshade spp.
Black medic	Field violet	PA smartweed
Carpetweed	FL beggarweed	Persian speedwell
Catchweed bedstraw	FL pusley	Pigweed spp.
Coast fiddleneck	Hairy beggarticks	Prostrate knotweed
Common dayflower	Hemp sesbania	Sheperd's-purse
Common lambsquarters	Henbit	Smallflowered bittercress
Common mallow	Ivyleaf speedwell	Wild buckwheat
Common purslane	Kochia	Wild lettuce
Corn speedwell	Ladysthumb	Wild mustard
Deadnettle spp.	Little mallow	

Category 2: Used in combinations with phenoxy-type postemergence broadleaf herbicides

Apply RightLine ADDIT Herbicide at rates of 0.9 to 1.5 fluid ounces of product per acre (0.021 to 0.034 fluid ounces per 1000 square feet) in tank-mix combinations with amines, esters, and salts of 2,4-D, dichlorprop, dicamba, mecoprop, MCPA and various combinations of these products for control of seedling and mature weeds listed in Category 1 plus the following:

CA burclover	FL betony	Prickly sida
Canada thistle	Ground ivy	Red sorrel
Carolina geranium	Healall	Redstem filaree
Common chickweed	Henbit	Spotted spurge
Common groundsel	Horseweed (Conyza Canadensis)	VA buttonweed
Common ragweed	Lawn burweed	White clover
Cutleaf eveningprimrose	Old-world diamond-flower	Wild violet
Dandelion	Parsley-piert	Yellow wood sorrel
Dollarweed	Perennial sowthistle	
Fleabane	Plantain spp.	

(continued)

Category 3: Used in combinations with other preemergence and postemergence herbicides registered for use in ornamental turf

RightLine ADDIT Herbicide may be applied at rates of 0.9 to 1.7 fluid ounces of product per acre (0.02 to 0.04 fluid ounces per 1000 square feet) in tank-mix combinations with clopyralid, foramsulfuron, glufosinate, glyphosate, halosulfuron-methyl, imazaquin, metsulfuron, and triclopyr for control of seedling annuals listed under Category 1 and for temporary burndown of weeds listed in Category 2. RightLine ADDIT Herbicide may be applied at rates of 0.9 to 1.7 fluid ounces of product per acre (0.02 to 0.04 fluid ounces per 1000 square feet) in a tank-mix combination with atrazine applied at 1.0 pounds of active ingredient per acre. Residual, long-term control of target weeds is as defined by the labeling of the companion product. For tank mixing with herbicides not listed above, follow the tank mix compatibility instructions under "Tank Mix Compatibility".

Category 4: Suppression

RightLine ADDIT Herbicide may be applied at rates of 1.0 to 2.1 fluid ounces of product per acre (0.023 to 0.048 fluid ounces per 1000 square feet) for temporary burndown of weeds listed in Category 2. RightLine ADDIT Herbicide may also be used for suppression and temporary burndown of algae (cyanobacteria) occurring in ornamental turf grasses and other landscape environments. Avoid spray drift onto desirable non-target susceptible plants such as vegetables, flowers, ornamentals, trees, shrubs.

RightLine ADDIT Herbicide Use Rate Conversion Table

fl. oz. product/acre	fl. oz. product/1,000 sq. ft.	ml product/1,000 sq. ft.	Lb. AI/Acre
0.9	0.021	0.6	0.013
1	0.023	0.7	0.015
1.2	0.028	0.8	0.018
1.3	0.030	0.9	0.019
1.5	0.034	1.0	0.022
1.6	0.037	1.1	0.024
1.7	0.039	1.2	0.025
1.8	0.041	1.2	0.027
1.9	0.044	1.3	0.028
2.1	0.048	1.4	0.031

(continued)

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide storage

DO NOT use or store in or around the home.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. **DO NOT** put formulated or dilute material into food or drink containers. **DO NOT** contaminate other pesticides, fertilizers, water, food, or feed by inappropriate storage or disposal.

To confine spill: Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged container in a larger holding container. Identify contents per required hazardous waste labeling regulations.

Pesticide Disposal

Waste resulting from the use of this product may be disposed of at an approved waste disposal facility.

Container Handling

NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS): DO NOT reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES and LIMITATIONS OF LIABILITY.** **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of RightLine, LLC. All such risks shall be assumed by the user or buyer. **DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, RightLine, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of RightLine, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, RightLine, LLC disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at RightLine, LLC's election, the replacement of product.