

FUNGICIDE

AZOXY

2SC SELECT™

LONG-TERM PROTECTION IN A CONVENIENT LIQUID FORMULATION.



Azoxy 2SC Select™ is the equivalent liquid formulation to the popular WDG azoxystrobin products. The suspended concentrate formulation allows for convenient tank mixing and application. The highly systemic formula in **Azoxy 2SC Select** delivers plant protection through both foliar and root absorption.

PRODUCT FEATURES

- Protection to the whole plant for up to 28 days.
- Convenient liquid formulation for easy tank mixing.
- Defends against both soil-borne and foliar diseases.
- Complete root-to-leaf distribution.
- Constant delivery for even distribution to new growth.
- Greater degree of root uptake when compared to other strobilurins.
- Prevents spore germination, thereby preventing many fungal infections.
- Does not impact honey bees and other beneficials.

LABELED USES

- Golf turf
- Residential turf
- Commercial turf
- Parks/Rec
- Athletic fields

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT

22.9% azoxystrobin

FRAC GROUP

11

FORMULATION

Suspension concentrate

CHEMICAL FAMILY

Strobilurin/QoI inhibitors

COMPARE TO

Heritage® TL

MODE OF ACTION

Acropetal penetrant

RESTRICTED USE

No

SIGNAL WORD

Caution

PACKAGING

12 x 1 pt, 12 x 1 qt, 4 x 1 gal, 2 x 2.5 gal

EPA REG NUMBER

89442-21

AZOXY 2SC SELECT™

LONG-TERM PROTECTION IN A CONVENIENT LIQUID FORMULATION.

APPLICATION INFORMATION

Azoxy 2SC Select™ may be applied as a foliar spray, in alternating spray programs, or in tank mixes with other registered pesticides.

Azoxy 2SC Select is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides, and biological control products. If tank mixes are desired, observe all directions, precautions, and limitations on labeling of all products used.

Azoxy 2SC Select is recommended for control of anthracnose, brown patch, cool weather brown patch (yellow patch), *Fusarium* patch, gray leaf spot, gray snow mold (*Typhula* blight), leafspot, melting out, necrotic ring spot, pink patch, pink snow mold, *Pythium* blight, *Pythium* root rot, red thread, *Rhizoctonia* large patch, southern blight, spring dead spot, summer patch, take-all patch, and *Zoysia* patch on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, and athletic fields.

Azoxy 2SC Select is most effective when applied prior to disease development. Mix **Azoxy 2SC Select** with the required amount of water and apply as a dilute spray application in 2-4 gallons of water per 1000 square feet (87-174 gallons per acre). Repeat applications at specified intervals for as long as required. For spot treatments, use 0.4 oz. **Azoxy 2SC Select** per 1 to 2 gallons of water. **DO NOT** apply more than 9.6 quarts product/acre/year (7.1 fl. oz. product/1000 square feet/year). Make applications by ground only.

LABELED DISEASES

Anthracnose	Gray leaf spot	Pink patch	<i>Rhizoctonia</i> large patch	<i>Typhula</i> blight
Brown patch	Gray snow mold	Pink snow mold	Spring dead spot	Yellow patch
Cool weather brown patch	Leaf spot	<i>Pythium</i> blight	Southern blight	<i>Zoysia</i> patch
<i>Fusarium</i> patch	Melting out	<i>Pythium</i> root rot	Summer patch	
	Necrotic ring spot	Red thread	Take-all patch	



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. Albaugh and the A logo are registered trademarks and Azoxy 2SC Select is a trademark of Albaugh, LLC. Heritage is a registered trademark of Syngenta Group Company. Always read and follow label restrictions, precautions, and directions for use. This product may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.